Approximate black holes from a variable cosmological constant

Maurice H. P. M. van Putten
Massachusetts Institute of Technology, Cambridge, MA 02139

arXiv:gr-qc/9802004v2 27 Apr 1998

The small or zero cosmological constant, Λ, probably results from a macroscopic cancellation mechanism of the zero-point energies. However, nearby horizon surfaces any macroscopic mechanism is expected to result in imperfect cancellations. A phenomenological description is given for the residual variable cosmological constant. In the static, spherically symmetric case it produces approximate black holes. The model describes the case of exponential decay by $\Box \ln \Lambda = -3a$, were a is a positive constant.

PACS numbers: 04.70.-s

The modern saga of the cosmological constant problem concerns the divergence of the zero-point energies [1]. A particularly compelling mechanism for cancellation of the resulting cosmological constant has been proposed by Coleman in the Euclidean path integral approach [2]. Coleman's analysis predicts a zero expectation value of the cosmological constant by integration over thin but extended wormholes. However, a horizon surface is expected to perturb any such macroscopic cancellation mechanism, by suppression of those wormholes which would tend to penetrate the surface. This perturbation suggests the possibility for a finite cosmological constant to reappear in the vicinity of horizon surfaces. In this *Letter*, a simple phenomenological approach is described to illustrate this scenario, using a self-consistent classical description of a variable cosmological constant.

A space-time with finite cosmological constant, Λ, filled with a fluid with stress-energy tensor, $T_{\alpha\beta}$, gives rise to the equations of motion

$$G_{\alpha\beta} = -\Lambda g_{\alpha\beta} + S_{\alpha\beta} \equiv T_{\alpha\beta}. \tag{1}$$

We shall look for time-independent, spherically symmetric solutions in the Schwarzschild line-element

$$\mathrm{d}s^2 = e^{\lambda}\mathrm{d}r^2 + r^2(\mathrm{d}\theta^2 + \sin^2\theta \mathrm{d}\phi^2) - e^{\nu}\mathrm{d}t^2. \tag{2}$$

A spherical region with finite Λ must be suitably closed by a shell with positive hoop stress to sustain the negative (radial) pressure inside. The shell also resides in a deep gravitational potential well, and therefore is expected to accumulate radiation and particles. If the particles interact infrequently (relative to the orbital time-scale), the stress tensor will develop an anisotropic pressure dominated by random orbital motion. In the limit of zero radial pressure of the fluid, its stress-energy tensor becomes

$$S_{\alpha\beta} = (E+P)u_\alpha u_\beta + P k_{\alpha\beta}, \tag{3}$$

where $u^\beta = (0,0,0,e^{-\nu/2})$ is the the average velocity four-vector of the particles,

$$k_{\alpha\beta} = g_{\alpha\beta} - (\partial^\gamma\Lambda\partial_\gamma\Lambda)^{-1}\partial_\alpha\Lambda\partial_\beta\Lambda \tag{4}$$

is a projection tensor, E is the energy density of the fluid and P is its two-dimensional pressure. Hence, we have

$$T_\alpha^\beta = -\mathrm{diag}(\Lambda, M, M, \Lambda + E), \tag{5}$$

were we have set $M = \Lambda - P$. It may be seen that the quantities Λ, E and P are scalars. The equations of energy-momentum conservation,

$$\nabla_\alpha T^\alpha_\beta = \partial_\alpha T^\alpha_\beta + \Gamma^\alpha_{\alpha\gamma}T^\gamma_\beta - \Gamma^\gamma_{\alpha\beta}T^\alpha_\gamma = 0, \tag{6}$$

reduce to

$$-\Lambda' + \Gamma^\alpha_{\alpha r}T^r_r - \Sigma_\alpha \Gamma^\alpha_{\alpha r}T^\alpha_\alpha = 0. \tag{7}$$

With the tabulated expressions for the connections (e.g.: [3]), this obtains

$$\Lambda' = \frac{2}{r}(M - \Lambda) + \frac{\nu'}{2}E. \tag{8}$$

At this point, recall the the weak energy condition on the stress-energy tensor

$$\inf T_{\alpha\beta}\xi^\alpha\xi^\beta = \Lambda + E \geq 0, \tag{9}$$

were the infimum is taken over all time-like vectors ξ^β. Clearly, (9) allows for the fluid to consist not only of particles ($E_p > 0$), but also of negative mass holes ($E_h < 0$) in regions with $\Lambda > 0$. Thus, coexistence of particles and holes makes it possible to have positive hoop stress, even with $E = 0$.

We will look for Schwarzschild solutions which join to space-time with a variable cosmological constant in the neighborhood of the origin, that is

$$e^\lambda \sim \left(1 - \tfrac{\Lambda_0 r^2}{3}\right)^{-1},\ e^\nu \sim 1 - \tfrac{\Lambda_0 r^2}{3},\ E, P \sim 0, \tag{10}$$

were Λ_0 is the value of the cosmological constant at the origin. The relevant Einstein equations are those for G^r_r and G^t_t,

$$\begin{aligned} e^{-\lambda}(\tfrac{\nu'}{r}+\tfrac{1}{r^2})-\tfrac{1}{r^2} =& -\Lambda, \\ -e^{-\lambda}(\tfrac{\lambda'}{r}-\tfrac{1}{r^2})-\tfrac{1}{r^2} =& -(\Lambda+E). \end{aligned} \tag{11}$$

These may be written as

$$\begin{aligned} r\nu' =& -1+e^{\lambda}(1-r^2\Lambda), \\ r\lambda' =& \ 1-e^{\lambda}(1-r^2[\Lambda+E]). \end{aligned} \tag{12}$$

Whenever $\Lambda = E = P = 0$ beyond some $r = r_0$, (12) recovers the Schwarzschild solution, though perhaps after a rescaling of the time coordinate (in case of $\lambda+\nu \neq 0$) to maintain the familiar relationship $-g_{tt}g_{rr} = 1$. Clearly, the simplest case follows with $E = 0$, which is allowed in our particle-hole model if any gravitational separation of particles and holes can be neglected on the scale of variation in Λ.

The simplest case, therefore, is given by a constitutive relation only for the anisotropy, P, in the pressure. The regularity condition of P at the origin, $P = O(r^2)$ for small r, translates covariantly to

$$P = \frac{\partial^\gamma\Lambda\partial_\gamma\Lambda}{F(\lambda)}, \tag{13}$$

were $F(\Lambda)$ is a continuous function of Λ. For example, $F(\lambda) = 2a\Lambda^\alpha$, so that

$$P = \frac{e^{-\lambda}}{2a\Lambda^\alpha}(\Lambda')^2 \tag{14}$$

giving a reduction of (8) to

$$\Lambda' = -are^{\lambda}\Lambda^\alpha. \tag{15}$$

Equation (15) shows a positioning of a transition layer in Λ at large e^λ, *i.e.*, at a surface of large red shift. In the case of $\alpha = 1$, the covariant generalization of (15) is

$$\Box \ln\Lambda = -3a. \tag{16}$$

A solution is illustrated in Figure 1. With $0 < \alpha < 1$, the distinguishing feature arises of Λ vanishing at finite r, as illustrated in Figure 2. The covariant generalization of (13) for $\alpha \neq 1$ is given by

$$\Box\Lambda^{(1-\alpha)} = -3a(1-\alpha); \tag{17}$$

for $\alpha > 1$, $\Lambda(r)$ decays algebraically in the far field limit.



Figure 1. Distributions of the metric quantities λ, ν and Λ as a function of radius in the case of $F = 2a\Lambda$ ($\alpha = 1$) for $a = 0.5, 1$ and $\Lambda = 5$ at the origin. A high red shift accompanies the transition layer in Λ. Outside of this layer, the cosmological constant is exponentially small, and space-time is essentially described by the exterior Schwarzschild solution. Inside this layer, the cosmological constant is effectively constant. The origin is singulaty free.



Figure 2. Distributions of the metric quantities λ, ν and Λ as a function of radius in the case of $F = 2a\Lambda^{0.5}$ ($\alpha = 0.5$) for $a = 1, 2$ and $\Lambda = 5$ at the origin. The transition layer is sharper compared with the previous case $F = 2a\Lambda$ with the same $a = 1$ as displayed in Figure 1. Furthermore, a bifurcation surface appears beyond which Λ vanishes identically, slightly outside the surface of high red shift. The origin is singularity free.

It is an open question whether (17) preserves non-negativity of $\Lambda^{(1-\alpha)}$ ($\Lambda_0 > 0$) in dynamical space-times. It may take numerical simulations to study the behavior of the surface of bifurcation (at which Λ becomes zero) in this setting. For this reason, a preliminary preference may be given to (16) which has $\alpha = 1$. It should be metioned that any terms on the right hand-side of (16-17) which vanish for $E = 0$ have been suppressed.

The existence of solutions can be seen from boundedness of λ. (12) shows that λ can blow up at a finite radius only when $r^2\Lambda > 1$. By (12), $r\lambda' \geq -(e^\lambda - 1)$, so that $\lambda \geq 0$. Then $\lambda' \leq re^\lambda\Lambda$ by the same equation. By (15), $re^\lambda\Lambda$ can be expressed in terms of Λ', so that

$$\lambda' \leq -\frac{1}{a}\frac{\Lambda'}{\Lambda^{\alpha-1}}. \tag{18}$$

It follows that

$$\lambda(r) \leq \lambda(r_0) + \frac{1}{a}\int_{\Lambda(r)}^{\Lambda(r_0)} \frac{d\Lambda}{\Lambda^{\alpha-1}}. \tag{19}$$

Clearly, if $0 < \alpha < 2$, we may take $r_0 = 0$, so that

$$0 \leq \lambda \leq \frac{1}{a(2-\alpha)}\Lambda_0^{2-\alpha} < \infty. \tag{20}$$

For $\alpha \geq 2$, choose r_0 such that $\Lambda(r_0)r_0^2 \geq 1$. For those $r \geq r_0$, where $\Lambda(r)r^2 \geq 1$, we have

$$\begin{array}{rcl} \lambda(r) & \leq & \lambda(r_0) + \frac{1}{a}\int_{1/r^2}^{\Lambda(r_0)} \Lambda^{1-\alpha} d\Lambda \\ & \leq & \lambda(r_0) + \frac{1}{a}\left\{ \begin{array}{ll} \ln \Lambda(r_0) r^2 & (\alpha = 2), \\ r^{2(\alpha-2)} & (\alpha > 2). \end{array} \right. \end{array} \tag{21}$$

Again, no blow-up at finite radius occurs.

The fully covariant stress-energy tensor is

$$S_{\alpha\beta} = E u_\alpha u_\beta + P h_{\alpha\beta} - \frac{\partial_\alpha \Lambda \partial_\beta \Lambda}{F(\Lambda)} \tag{22}$$

with $h_{\alpha\beta} = g_{\alpha\beta} + u_\alpha u_\beta$. The energy E has been included for consistency in view of the normalization $u^2 = -1$, so that the four equations of energy-momentum conservation,

$$\nabla_\alpha T^\alpha_\beta = 0 \tag{23}$$

can be satisfied with the three degrees of freedom in u^β plus the one degree of freedom in E. As we have seen, E will depart from zero only in genuinely dynamical space-times. The evolution equation for E and u^β satisfies

$$\nabla_\alpha (E u^\alpha u_\beta) + \nabla_\alpha S^\alpha_\beta - \partial_\beta \Lambda = 0; \tag{24}$$

in particular,

$$\nabla_\alpha (E u^\alpha) = u^\beta \partial_\beta \Lambda + u^\beta \nabla_\alpha S^\alpha_\beta \tag{25}$$

expresses the evolution of E.

The present solutions feature surfaces of arbitrarily high red shift, outside of which space-time is essentially that of general relativity. These solutions are related to the earlier solution of van Putten [7], which were proposed as as approximate black holes for numerical regularization. Spherically symmetric static solutions with surfaces of arbitrarily high red shift have recently also been described by ‘t Hooft [8,9], in an entirely different context. These solutions are both hampered by naked singularities at the origin (though of different kinds). Numerical simulations by Coptuik *et al*. [6] also show the solution of [7] to be unstable. This naturally raises the same question of stability of the present solutions, something which falls outside the scope of the present work. It should be emphasized that the present solutions are an improvement (over forementioned related solutions) in that these are free of any singularities.

In regards to the application to numerical relativity, note that the stress-energy tensor remains differentiable everywhere, though with a discontinuity in its first derivative if $0 < \alpha < 1$. This still provides a proper source term in the covariant, strictly hyperbolic formulation in terms of nonlinear wave equations by van Putten & Eardley [4], given by

$$\hat{\Box} \omega_{\alpha ab} - R^\gamma_\alpha \omega_{\gamma ab} - [\omega^\gamma, \nabla_\alpha \omega_\gamma]_{ab} = 16\pi \tau_{\alpha ab}, \tag{26}$$

were $\omega_{\alpha ab}$ are the tetrad connections of the tetrad elements $(e_a)_\alpha$ (here, Latin indices refer to the tetrad indices and Greek indices refer to the coordinates). The source term $\tau_{\alpha ab} = \tau_{\alpha\beta\gamma}(e_a)^\beta (e_b)^\gamma$, is given by

$$\tau_{\alpha\beta\gamma} = \nabla_{[\beta} T_{\gamma]\alpha} - \frac{1}{2} g_{\alpha[\gamma} \nabla_{\beta]} T, \tag{27}$$

were $T = T^\gamma_\gamma$ is the trace of the stress-energy tensor. The equations in the exterior vacuum, therefore, are

$$\hat{\Box} \omega_{\alpha ab} - [\omega^\gamma, \nabla_\alpha \omega_\gamma]_{ab} = 0. \tag{28}$$

These have been given a second-order numerical implementation in one dimension [5]. In the case of $0 < \alpha < 1$, $\tau_{\alpha ab}$ in (26) is at most discontinuous, due to second derivatives of Λ; for $\alpha \geq 1$, there are no discontinuities. It is expected, therefore, that for all $\alpha > 0$ the stress-energy tensor (22), plus $-\Lambda g_{\alpha\beta}$ and constitutive relation (13), can indeed be integrated into this approach of numerical relativity.

ACKNOWLEDGEMENTS

The author gratefully acknowledges stimulating discussions with Ira Wasserman in the very early stages of this work, Thomas Witelski, Hung Cheng, Daniel Freedman and Alan Guth; and S. Caveny for a careful reading. This work has received partial support from the MIT Sloan/Cabot Fund.

---

[1] S. Weinberg. *Rev. Mod. Phys.*, 61:1, 1989.
[2] S. Coleman. *Nucl. Phys. B*, 307:867, 1988.
[3] H. Stephani. *General Relativity* (Cambridge, 1990).
[4] M.H.P.M. van Putten and D.M. Eardley. *Phys. Rev. D*, 53:3056, 1996.
[5] M.H.P.M. van Putten. *Phys. Rev. D*, 55:4705, 1996.
[6] M.W. Choptuik, E.W. Hirschmann and S.L. Liebling. *Phys. Rev. D*, 55:6014, 1997.
[7] M.H.P.M. van Putten. *Phys. Rev. D*, 54:R5931, 1996.
[8] G. ’t Hooft. *gr-qc*/9706058, 1997.
[9] G. ’t Hooft. *gr-qc*/9711055, 1997.